


IN THE CLAIMS

1. (Original): A compound of the formula



wherein the bond of atoms C₂₂ and C₂₁ is a single or double bond;

m is 0 or 1;

n is 0, 1 or 2;

p is 0 or 1;

R₁ is C₁-C₁₂-alkyl, C₃-C₈-cycloalkyl or C₂-C₁₂-alkenyl;

R₂ is H, C₁-C₁₂-alkyl, C₁-C₁₂-haloalkyl, C₁-C₁₂-hydroxyalkyl, OH, halogen, -N₃, SCN, NO₂, CN, C₃-C₈-cycloalkyl unsubstituted or substituted by from one to three methyl groups, C₃-C₈-halocycloalkyl, C₁-C₁₂-alkoxy, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₁-C₆-alkoxy-C₁-C₆-alkoxy, C₁-C₆-alkoxy-C₁-C₆-alkoxy-C₁-C₆-alkyl, C₂-C₁₂-alkenyl, C₂-C₁₂-haloalkenyl, C₂-C₁₂-haloalkenyloxy, C₂-C₁₂-alkynyl, C₂-C₁₂-haloalkynyl, C₃-C₁₂-alkynyl-oxy, C₃-C₁₂-haloalkynyloxy, -P(=O)(OC₁-C₆-alkyl)₂, -Si(C₁-C₆-alkyl)₃, -(CH₂)-Si(C₁-C₆-alkyl)₃, -Si(OC₁-C₆-alkyl)₃, -N(R₉)₂, -(CH₂)-N(R₉)₂, wherein the two substituents R₉ are independent of each other, -C(=X)-R₇, -(CH₂)-C(=X)-R₇, -O-C(=X)-R₇, -(CH₂)-O-C(=X)-R₇, -S-C(=X)-R₇, -(CH₂)-S-C(=X)-R₇, -NR₉C(=X)R₇, -(CH₂)-NR₉C(=X)R₇, -NR₉NHC(=X)-R₇, -NR₉-OR₁₀, -(CH₂)-NR₉-OR₁₀, -SR₉, -S(=O)R₁₁, -S(=O)₂R₁₁, aryl, heterocyclyl, aryloxy or heterocyclyloxy; wherein the aryl, heterocyclyl, aryloxy and heterocyclyloxy radicals are unsubstituted or, depending upon the possibilities of substitution at the ring, mono- to penta-substituted by substituents selected from the group consisting of OH, halogen, CN, NO₂, SCN, -N₃, C₁-C₁₂-alkyl, C₃-C₈-cycloalkyl, C₁-C₁₂-haloalkyl, C₁-C₁₂-alkoxy, C₁-C₁₂-haloalkoxy, C₁-C₁₂-alkylthio, C₁-C₁₂-haloalkylthio, C₁-C₆-alkoxy-C₁-C₆-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, C₂-C₁₂-haloalkenyl, C₂-C₁₂-haloalkenyloxy, C₂-C₁₂-haloalkynyl, C₃-C₁₂-alkynyloxy, C₃-C₁₂-haloalkynyloxy and phenoxy;

or, when p is 1, R₂ together with R₃ is a bond;

or R₂ together with R₄ is =O or =S;

or R₂ together with R₄ form with the carbon to which they are bound a three- to seven-membered ring, which may be monocyclic or bicyclic, and may be saturated or unsaturated, and that may contain one or two hetero atoms selected from the group consisting of N, O and S, and which is either unsubstituted or independently of one another mono- to pentasubstituted with substituents selected from OH, =O, SH, =S, halogen, CN, -N₃, SCN, NO₂, aryl, C₁-C₁₂alkyl, C₃-C₈cycloalkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₁-C₁₂haloalkylthio, C₁-C₆alkoxy-C₁-C₆alkyl, C₂-C₈alkenyl, C₂-C₈alkynyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂haloalkynyl, C₃-C₁₂alkynyloxy, C₃-C₁₂haloalkynyloxy, phenoxy, phenyl-C₁-C₆alkyl, -N(R₉)₂ wherein the two R₉ are independent of each other, C₁-C₆alkylsulfinyl, C₃-C₈cycloalkylsulfinyl, C₁-C₆haloalkylsulfinyl, C₃-C₈halocycloalkylsulfinyl, C₁-C₆alkylsulfonyl, C₃-C₈cycloalkylsulfonyl, C₁-C₆haloalkylsulfonyl and C₃-C₈halocycloalkylsulfonyl; or

R₂ together with R₄ is =NN(R₁₂)₂, wherein the two substituents R₉ are independent of each other;

or, when p is 0, R₂ together with R₄ and R₆ is =N;

or when p is 0, R₂ together with R₆ is -NOR₁₂ or =NN(R₁₂)₂, wherein the two substituents R₉ are independent of each other;

R₃ is H, C₁-C₁₂-alkyl, halogen, halo-C₁-C₂alkyl, CN, -N₃, SCN, NO₂, C₃-C₈cycloalkyl unsubstituted or substituted by from one to three methyl groups, C₃-C₈halocycloalkyl, C₁-C₁₂alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₆alkoxy-C₁-C₆alkoxy-C₁-C₆alkyl, C₃-C₈cycloalkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₃-C₈cycloalkylthio, C₁-C₁₂haloalkylthio, C₁-C₁₂alkylsulfinyl, C₃-C₈cycloalkylsulfinyl, C₁-C₆haloalkylsulfinyl, C₃-C₈halocycloalkylsulfinyl, C₁-C₁₂alkylsulfonyl, C₃-C₈cycloalkylsulfonyl, C₁-C₁₂haloalkylsulfonyl, C₃-C₈halocycloalkylsulfonyl, C₂-C₈alkenyl, C₂-C₈alkynyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyloxy, -N(R₉)₂, wherein the two substituents R₉ are independent of each other, aryl, heterocyclyl, aryloxy or heterocyclyloxy; wherein the aryl, heterocyclyl, aryloxy and heterocyclyloxy radicals are unsubstituted or, depending upon the possibilities of substitution at the ring, mono- to penta-substituted by substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₁₂alkyl, C₃-C₈cycloalkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₁-C₁₂haloalkylthio, C₁-C₆alkoxy-C₁-C₆alkyl, C₂-C₈alkenyl, C₂-C₈alkynyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂haloalkynyl and C₃-C₁₂haloalkynyloxy;

or when p is 1, R₃ together with R₇ is a bond;

R₄ is H, C₁-C₁₂-alkyl, C₁-C₁₂-haloalkyl, C₁-C₁₂-hydroxyalkyl, OH, halogen, NO₂, CN, C₃-C₈cycloalkyl unsubstituted or substituted by from one to three methyl groups, C₃-C₈halocycloalkyl, C₁-C₁₂alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₆alkoxy-C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkoxy-C₁-C₆alkyl, C₂-C₁₂alkenyl,

C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂alkynyl, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyl-oxy, -P(=O)(OC₁-C₆alkyl)₂, -Si(C₁-C₆alkyl)₃, -(CH₂)-Si(C₁-C₆alkyl)₃, -Si(OC₁-C₆alkyl)₃, -N(R₉)₂, -(CH₂)-N(R₉)₂, wherein the two substituents R₉ are independent of each other, -C(=X)-R₇, -(CH₂)-C(=X)-R₇, -O-C(=X)-R₇, -(CH₂)-O-C(=X)-R₇, -S-C(=X)-R₇, -(CH₂)-S-C(=X)-R₇, -NR₉C(=X)R₇, -(CH₂)-NR₉C(=X)R₇, -NR₉NHC(=X)-R₇, -NR₉-OR₁₀, -(CH₂)-NR₉-OR₁₀, -SR₉, -S(=O)R₁₁, -S(=O)₂R₁₁, aryl, heterocyclyl, aryloxy or heterocyclyloxy; wherein the aryl, heterocyclyl, aryloxy and heterocyclyloxy radicals are unsubstituted or, depending upon the possibilities of substitution at the ring, mono- to penta-substituted by substituents selected from the group consisting of OH, halogen, CN, NO₂, C₁-C₁₂alkyl, C₃-C₈cycloalkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₁-C₁₂haloalkylthio, C₁-C₆alkoxy-C₁-C₆alkyl, C₂-C₈alkenyl, C₂-C₈alkynyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyloxy and phenoxy;

or R₄ together with R₂ forms =O or -S;

or when p is 1, R₄ together with R₅ is a bond;

or, when p is 0, together with R₂ and R₆ is ≡N;

R₅ and R₆ independently of each other are H, C₁-C₁₂-alkyl, -N₃, CN, NO₂, OH, SH, halogen, halo-C₁-C₂alkyl, hydroxy-C₁-C₂alkyl, C₃-C₈cycloalkyl that is unsubstituted or substituted by from one to two methyl groups, C₃-C₈halocycloalkyl, C₁-C₁₂alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₆alkoxy-C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₈cycloalkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂haloalkylthio, C₂-C₈alkenyl, C₂-C₈alkynyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyloxy, -P(=O)(OC₁-C₆alkyl)₂, -CH₂-P(=O)(OC₁-C₆alkyl)₂, -Si(OC₁-C₆alkyl)₃, -N(R₉)₂, -O-N(R₉)₂, wherein the two substituents R₉ are independent of each other, -C(=X)-R₇, -CH=NOH, -CH=NOC₁-C₆alkyl, -O-C(=X)-R₇, -S-C(=X)-R₇, -NR₉C(=X)R₇, -NR₉NHC(=X)-R₇, -NR₉-OR₁₀, -SR₉, -S(=O)R₁₁, -S(=O)₂R₁₁, -CH₂-S(=O)₂R₁₁, aryl, aryloxy, benzyloxy, -NR₉-aryl, heterocyclyl, heterocyclyloxy, -NR₉-heterocyclyl, -CH₂-aryl, -CH₂-O-aryl, -CH₂-NR₉-aryl, -CH₂-NR₉-C₁-C₂alkyl, -CH₂-heterocyclyl, -CH₂-O-heterocyclyl and -CH₂-NR₉-heterocyclyl; wherein the aryl, aryloxy, benzyloxy, -NR₉-aryl, heterocyclyl, heterocyclyloxy and -NR₉-heterocyclyl radicals are unsubstituted or, depending upon the possibilities of substitution at the ring, mono- to penta-substituted by substituents selected from the group consisting of OH, =O, SH, =S, halogen, CN, NO₂, C₁-C₁₂alkyl, C₃-C₈cycloalkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₁-C₁₂haloalkylthio, C₁-C₆alkoxy-C₁-C₆alkyl, C₂-C₈alkenyl, C₂-C₈alkynyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyloxy, phenoxy, methylenedioxy, NH₂, NH(C₁-C₁₂alkyl), N(C₁-C₁₂alkyl)₂ and C₁-C₆alkylsulfinyl; or

R₅ and R₆ are, together with the carbon atom to which they are bound, a five- to seven-membered ring, which may be saturated or unsaturated, and which may contain one or two members selected from the group consisting of O, NR₈ and S; and which is optionally substituted with one to three substituents selected from

C₁-C₁₂-alkyl, CN, NO₂, OH, halogen, halo-C₁-C₂alkyl, C₃-C₈cycloalkyl C₃-C₈halocycloalkyl, C₁-C₁₂alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₆alkoxy-C₁-C₆alkoxy-C₁-C₆alkyl, C₃-C₈cycloalkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₃-C₈cycloalkylthio, C₁-C₁₂haloalkylthio, C₂-C₁₂alkenyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂alkynyl, C₂-C₁₂haloalkynyl and C₃-C₁₂haloalkynyloxy;

or when p is 1, R₅ together with R₄ is a bond;

or, when p is 0, R₆ together with R₂ and R₄ is $\equiv N$;

R₇ is H, OH, C₁-C₁₂alkyl, C₁-C₁₂haloalkyl, C₂-C₁₂alkenyl, C₂-C₁₂alkynyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyloxy, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₆-alkoxy-C₁-C₆alkyl, C₁-C₆alkoxy-C₁-C₆alkoxy, C₂-C₈alkenyloxy, C₁-C₈alkynyloxy, -N(R₈)₂ wherein the two R₈ are independent of each other, aryl, aryloxy, benzyloxy, heterocyclyl, heterocyclyloxy or heterocyclylmethoxy; and wherein the aryl, aryloxy, benzyloxy, heterocyclyl and heterocyclyloxy radicals are unsubstituted or, depending upon the possibilities of substitution at the ring, mono- to penta-substituted by substituents selected from the group consisting of halogen, CN, NO₂, C₁-C₁₂alkyl, C₃-C₈cycloalkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₁-C₁₂haloalkylthio, C₁-C₆alkoxy-C₁-C₆alkyl, C₂-C₈alkenyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₈alkynyl, C₂-C₁₂haloalkynyl and C₃-C₁₂haloalkynyloxy;

R₈ is H, C₁-C₆alkyl that is optionally substituted with one to five substituents selected from the group consisting of halogen, C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkoxy, C₂-C₁₂alkenyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂alkynyl, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyloxy, hydroxy and cyano, C₃-C₈-cycloalkyl, aryl, benzyl or heteroaryl; wherein the aryl, benzyl and heteroaryl radicals are unsubstituted or, depending on the possibilities of substitution on the ring, mono- to trisubstituted by substituents selected from the group consisting of OH, halogen, CN, NO₂, C₁-C₁₂alkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₂-C₁₂alkenyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂alkynyl, C₂-C₁₂haloalkynyl, C₃-C₁₂haloalkynyloxy and C₁-C₁₂haloalkylthio;

R₉ is H, C₁-C₆alkyl, C₁-C₆cycloalkyl, C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₆alkoxy-C₁-C₆alkoxy-C₁-C₆alkyl, C₂-C₁₂alkenyl, C₂-C₁₂alkynyl, benzyl, aryl or heteroaryl;

R₁₀ H, C₁-C₆alkyl that is optionally substituted with one to five substituents selected from the group consisting of halogen, C₁-C₆alkoxy, NO₂, hydroxy and cyano, C₁-C₁₂haloalkyl, C₂-C₁₂alkenyl, C₂-C₁₂haloalkynyl, C₂-C₁₂haloalkenyl, C₂-C₁₂alkynyl, C₃-C₈-cycloalkyl, aryl, benzyl or heteroaryl; wherein the aryl, benzyl and heteroaryl radicals are unsubstituted or, depending on the possibilities of substitution on the ring, mono- to trisubstituted by substituents selected from the group consisting of OH, halogen, CN, NO₂, C₁-C₁₂alkyl, C₁-C₁₂haloalkyl, C₁-C₁₂alkoxy, C₁-C₁₂haloalkoxy, C₁-C₁₂alkylthio, C₁-C₁₂haloalkylthio, C₂-C₁₂alkenyl, C₂-C₁₂haloalkenyl, C₂-C₁₂haloalkenyloxy, C₂-C₁₂alkynyl, C₃-C₁₂haloalkynyl and C₃-C₁₂haloalkynyloxy;

R_{11} is H, C_1 - C_6 alkyl that is optionally substituted with one to five substituents selected from the group consisting of halogen, C_1 - C_6 alkoxy, hydroxy and cyano, $-N(R_9)_2$ wherein the two substituents R_9 are independent of each other, C_3 - C_8 cycloalkyl, C_1 - C_8 halocycloalkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} haloalkenyl, C_2 - C_{12} haloalkenyloxy, C_2 - C_{12} alkynyl, C_3 - C_{12} haloalkynyl, C_3 - C_{12} haloalkynyloxy, aryl, benzyl or heteroaryl; wherein the aryl, benzyl and heteroaryl radicals are unsubstituted or, depending on the possibilities of substitution on the ring, mono- to trisubstituted by substituents selected from the group consisting of OH, halogen, CN, NO_2 , C_1 - C_{12} alkyl, C_1 - C_{12} haloalkyl, C_1 - C_{12} alkoxy, C_1 - C_{12} haloalkoxy, C_1 - C_{12} alkylthio, C_1 - C_{12} haloalkylthio, C_2 - C_{12} alkenyl, C_2 - C_{12} haloalkenyl, C_2 - C_{12} haloalkenyloxy, C_2 - C_{12} alkynyl, C_2 - C_{12} haloalkynyl and C_3 - C_{12} haloalkynyloxy;

R_{12} is H, C_1 - C_6 alkyl, C_1 - C_6 cycloalkyl, C_1 - C_6 alkoxy- C_1 - C_6 alkyl, C_1 - C_6 alkoxy- C_1 - C_6 alkoxy- C_1 - C_6 alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} alkynyl, $-C(=O)C_1$ - C_6 alkyl, $-C(=O)OC_1$ - C_6 alkyl, $-SO_2C_1$ - C_6 alkyl, benzyl, aryl, heteroaryl;

X is O or S;

or, if appropriate, an E/Z isomer, E/Z isomer mixture and/or tautomer thereof, in each case in free form or in salt form;

with the proviso, that the group $R_n-[C(R_1)(R_3)]_p-C(R_2)(R_4)-[CH_2]_n$, which is attached to the \square -position of the compound of the formula (I), is not $NC-CH_2-$ or $HOOC-CH_2-$ when m is 1 and the bond between atoms 22 and 23 is a single bond.

2. (Previously Presented): A pesticide composition which contains at least one compound of the formula (I) as described in claim 1 as active compound and at least one auxiliary.

3. (Previously Presented): A method for controlling pests comprising applying a composition as described in claim 2 to the pests or their habitat.

4. (Previously Presented): A process for preparing a composition as described in claim 2 comprising intimately mixing and/or grinding the active compound with at least one auxiliary.

5. (Cancelled).

6. (Cancelled).

7. (Previously Presented): A method for protecting plant propagation material, wherein the propagation material or the location where the propagation material is planted is treated, comprising applying a composition as described in claim 2.

8. (Currently Amended): Plant propagation material treated ~~in accordance with the method~~ composition described in claim 7 2.